arXiv:0804.3515v1 [astro-ph] 22 Apr 2008

Submitted to *The Astrophysical Journal Letters*

The Hard VHE γ-Ray Emission in High-Redshift TeV Blazars: Comptonization of Cosmic Microwave Background Radiation in an Extended Jet?

Markus Böttcher[1], Charles D. Dermer[2], and Justin D. Finke[2,3]

ABSTRACT

Observations of very-high-energy (VHE, $E > 250$ GeV) γ-ray emission from several blazars at $z > 0.1$ have placed stringent constraints on the elusive spectrum and intensity of the intergalactic infrared background radiation (IIBR). Correcting their observed VHE spectrum for $\gamma\gamma$ absorption even by the lowest plausible level of the IIBR provided evidence for a very hard (photon spectral index $\Gamma_{\rm ph} < 2$) intrinsic source spectrum out to TeV energies. Such a hard VHE γ-ray spectrum poses a serious challenge to the conventional synchrotron-self-Compton interpretation of the VHE emission of TeV blazars and suggests the emergence of a separate emission component beyond a few hundred GeV. Here we propose that such a very hard, slowly variable VHE emission component in TeV blazars may be produced via Compton upscattering of Cosmic Microwave Background (CMB) photons by shock-accelerated electrons in an extended jet. For the case of 1ES 1101-232, this component could dominate the bolometric luminosity of the extended jet if the magnetic fields are of the order of typical intergalactic magnetic fields ($B \sim 10\,\mu$G) and electrons are still being accelerated out to TeV energies ($\gamma \gtrsim 4 \times 10^6$) on kiloparsec scales along the jet.

Subject headings: galaxies: active — BL Lac Objects: individual (1ES 1101-232) — gamma-rays: theory — radiation mechanisms: non-thermal

---

[1] Astrophysical Institute, Department of Physics and Astronomy,
Clippinger 339, Ohio University, Athens, OH 45701; boettchm@ohio.edu

[2] Space Science Division, Code 7653, Naval Research Laboratory,
Washington, DC 20375-5352

[3] NRL/NRC Research Associate

1. Introduction

More than a dozen blazars have now been detected as sources of VHE γ-ray emission. Almost all of these objects are high-frequency-peaked BL Lac objects (HBLs) at low redshifts ($z \lesssim 0.2$). The low redshifts can be attributed to the absorption of VHE γ-rays by the IIBR. However, the level of the IIBR in the local universe and its evolution with redshift are very difficult to measure directly (e.g., Hauser & Dwek 2001). Consequently, the intensity and spectral energy distribution of the IIBR and the resulting optical depth of the Universe to $\gamma\gamma$ absorption and pair production are still a matter of intensive debate (see, e.g., Dwek & Krennrich 2005; Aharonian et al. 2006; Stecker et al. 2006; Stecker & Scully 2006; Mazin & Raue 2007; Aharonian et al. 2007a).

The $\gamma\gamma$ opacity of the Universe for GeV – TeV photons is generally increasing with increasing photon energy and with increasing redshift. Therefore, any observed VHE γ-ray spectrum with a local photon spectral index $\Gamma_{\rm obs}$ (for $\Phi_{\rm obs}(E)$ [ph/(cm^2 s)] $\propto E^{-\Gamma_{\rm obs}}$) corresponds to an intrinsic spectral index $\Gamma_{\rm int} < \Gamma_{\rm obs}$ at the source, i.e., the intrinsic VHE γ-ray emission at the source must not only be more luminous, but its spectrum must also be harder than the observed one. A strict upper limit on the intensity of the IIBR at the maximum of its spectral energy distribution (SED) around 1.5 μm of $\sim$ 14 nW m^{-2} sr^{-1} has been inferred by Aharonian et al. (2006) based on the argument that any plausible mechanism of particle acceleration and radiative cooling will result in an effective particle spectral lindex $p \geq 2$ (where $n_e(\gamma) \propto \gamma^{-p}$). For any plausible radiation mechanism (most notably Compton scattering), this would correspond to a limiting intrinsic photon spectral index of $\Gamma_{\rm int} \geq 1.5$. According to Aharonian et al. (2006) (see also, Mazin & Raue 2007) the upper limit on the 1.5 μm IIBR intensity from the requirement $\Gamma_{\rm int} \geq 1.5$ for the case of 1ES 1101-232 is now rather close to the lower limit from direct galaxy counts (e.g. Madau & Pozetti 2000). Requiring that $\Gamma_{\rm int} > 2$ (Dermer 2007) allows a low intensity IIBR that is compatible with observations (Hauser & Dwek 2001). However, recent numerical simulations by Stecker et al. (2007) indicated that particle spectra with $q < 1.5$ can result from diffusive shock acceleration at relativistic shocks, which loosens the constraints on the IIBR inferred by Aharonian et al. (2006). In this context, it should also be mentioned that acceleration at shear flows (e.g., Stawarz & Ostrowski 2002; Rieger & Duffy 2006) as well as second-order Fermi acceleration (e.g. Virtanen & Vainio 2005) may lead to ultrarelativistic particle acceleration into particle distributions as hard as $q \sim 1$ or even injection of particle distributions with low-energy cutoffs at ultrarelativistic energies (Derishev 2003; Stern & Poutanen 2008). Subsequent radiative cooling will then produce relativistic electron spectra with a canonical index of $p = 2$.

Based on the IIBR spectrum inferred from the constraints described above, Aharonian et al.

(2007a) reconstructed the intrinsic VHE γ-ray spectrum of the HBL 1ES 1101-232 at $z = 0.186$, which has recently been detected as a VHE γ-ray source by the H.E.S.S. array (Aharonian et al. 2006). Their reconstruction was based on the limiting value of the intrinsic spectral index of $\Gamma_{\rm int} \sim 1.5$, corresponding to $p \sim 2.0$ (for Compton emission in the Thomson limit). Even for the minimum plausible value of the IIBR intensity, they inferred that the peak of the high-energy component of the SED must be located at $E_p \geq 3$ TeV. While the broadband SEDs of HBLs detected at VHE γ-rays can generally be fit with basic synchrotron-self-Compton (SSC) models (see, however Finke et al. 2008, for PKS 2155-304), they are very hard-pressed to reproduce such an extremely hard GeV – TeV spectrum (see, e.g., Aharonian et al. 2007a). This is because at those energies the SSC spectrum exhibits strong intrinsic curvature due to Klein-Nishina effects: the number of soft photons that can efficiently be scattered (in the Thomson regime) steadily decreases with increasing scattered (γ-ray) photon energy. Therefore, even an electron spectrum with $p = 2$ will generally produce an SSC spectrum with local spectral index at GeV – TeV energies of $\Gamma_{\rm int} > 1.5$. One way to overcome this problem could be a very high low-energy cutoff of the electron energy distribution ($\gamma_{\rm min} \gtrsim 10^5$), as recently suggested by Katarzyński et al. (2006). However, maintaining such a high-energy cutoff over the relevant radiative time scale would require that the emission region is radiatively inefficient, which is unlikely.

The extremely hard intrinsic GeV – TeV spectrum of 1ES 1101-232 might suggest that there is an additional, very hard spectral component emerging above the SSC emission at those photon energies. Here we propose that such a component could be produced through Compton upscattering of the cosmic microwave background (CMB) radiation in the extended region of the jet. This mechanism has previously been considered as a way to explain the hard X-ray emission in the extended jets of several radio quasars and radio galaxies resolved with *Chandra* (e.g., Sambruna et al. 2004), see however Atoyan & Dermer (2004). The X-ray spectra of many of those extended jet components are very hard with $\Gamma_x \sim 1.5$, suggesting that even in the case of mildly beamed jets of non-blazar radio-loud AGN such emission might extend into the MeV – GeV range. It seems then worthwhile to investigate whether in the case of blazars, such an emission component could even extend into the GeV – TeV regime and produce a quasi-steady plateau of very hard VHE emission.

Throughout this paper, we refer to α as the energy spectral index, F_ν [Jy] $\propto \nu^{-\alpha}$ and the photon spectral index $\Gamma_{\rm ph} = \alpha + 1$. A cosmology with $\Omega_m = 0.3$, $\Omega_\Lambda = 0.7$, and $H_0 = 70$ km s^{-1} Mpc^{-1} is used. In this cosmology, and using the redshift of $z = 0.186$, the luminosity distance of 1ES 1101-232 is $d_L = 905$ Mpc.

2. Compton upscattering of the CMB to VHE γ-rays

The CMB at the source at redshift z is a thermal blackbody spectrum with a temperature $T_{\rm CMB} = (1+z) \times 2.72$ K. Its local energy density is $u_{\rm CMB} = (1+z)^4 \times 4.02 \times 10^{-13}$ ergs cm^{-3}. For the purpose of an analytical estimate, the CMB can be approximated as a monochromatic radiation field at a normalized mean photon energy $\epsilon_{\rm CMB} \equiv (h\nu_{\rm CMB}/m_e c^2) \approx (1+z) \times 1.2 \times 10^{-9}$. If the emission region is moving with respect to the local AGN frame with a bulk Lorentz factor Γ, the CMB radiation field is boosted into the comoving frame by a factor of Γ in photon energy and a factor of Γ^2 in energy density. In the following, we scale $\Gamma = 10\,\Gamma_1$, and, accordingly, the Doppler boosting factor $D \equiv (\Gamma[1 - \beta_\Gamma \cos\theta_{\rm obs}])^{-1} \equiv 10\,D_1$, where $\theta_{\rm obs}$ is the observing angle. Primed quantities refer to the co-moving frame of the emission region.

Efficient Compton upscattering of the CMB into the γ-ray regime with the observed very hard intrinsic spectrum of 1ES 1101-232, requires that the scattering occurs in the Thomson regime which is possible for electrons with Lorentz factors of $\gamma' \le (4\Gamma\epsilon'_{\rm CMB})^{-1} = 2.1 \cdot 10^7\,\Gamma_1^{-1}\,(1+z)^{-1}$. Such electrons will Compton scatter CMB photons up to energies of $\epsilon_{\rm s,max}^{\rm obs} \approx 5.3 \times 10^7\,D_1\,\Gamma_1^{-1}\,(1+z)^{-2}$ or $E_{\rm s,max}^{\rm obs} \approx 27\,D_1\,\Gamma_1^{-1}\,(1+z)^{-2}$ TeV in the observer's frame. Thus, if electrons can be accelerated to TeV energies, they will be able to upscatter CMB photons into the TeV regime in the Thomson limit. Specifically, to scatter a CMB photon up to 1 TeV in the observer's frame, a comoving electron Lorentz factor of $\gamma'_{\rm TeV} = 4.0 \times 10^6\,(D_1\Gamma_1)^{-1/2}$ is required. The radiative cooling time scale due to Compton cooling in the co-moving frame is

$$\tau'_{\rm CMB}(\gamma_{\rm TeV}) = 6.3 \times 10^3\,D_1^{1/2}\,\Gamma_1^{-3/2}\,(1+z)^{-1}\ {\rm yr}. \tag{1}$$

Based on this cooling time scale it will in principle be possible to accelerate electrons out to $\gamma'_{\rm TeV}$ for magnetic field values exceeding $B' \gtrsim 1.2 \times 10^{-12}\,D_1^{-1}\,\Gamma_1\,(1+z)^4$ G, which does not impose a serious constraint. CMB Compton cooling will dominate over synchrotron cooling for a co-moving magnetic field value of

$$B' \lesssim 3.2 \times 10^{-5}\,\Gamma_1\,(1+z)^2\ {\rm G}. \tag{2}$$

The resulting synchrotron emission produced by electrons with energy $\gamma'_{\rm TeV}$ will peak at $\nu_{\rm sy}^{\rm obs} \lesssim 2.19 \times 10^{16}\,(1+z)$ Hz, i.e., typically in the UV or soft X-ray bands. The dominance of CMB Compton cooling over synchrotron cooling is required in order for the CMB Compton emission to be at least at a comparable level to the synchrotron emission from the same electrons. The required low magnetic fields are much lower than the magnetic field strengths inferred for the presumably near-central (sub-pc) regions of the jet from where the bulk of the observed optical – X-ray – GeV γ-ray emission from blazars is generally believed to originate. This points towards an origin in the outer (pc – kpc scale) regions of the jet, where the magnetic fields might approach characteristic interstellar magnetic field values of a few tens of μG. The resulting magnetic jet luminosity will be $L_{\rm mag} \sim 1.5 \times 10^{39} \, R_{18}^2 \, \Gamma_1^4$ ergs s^{-1}, where R_{18} is the (poorly constrained) cross-sectional radius of the jet in units of 10^{18} cm, and thus several orders of magnitudes less than the total luminosity required to power the emission. This indicates that the jets must be particle dominated at the site of the VHE γ-ray production through CMB upscattering, even for a pc-scale transverse extent of the jet.

A key prediction from the inferred long cooling time scale given above is that the CMB Compton emission, as well as the synchrotron emission associated with the same emission region, will be slowly variable. This is another reason why the magnetic-field energy density in the VHE emitting region should be lower than the CMB energy density: otherwise this would contradict the observations of rapid variability throughout the synchrotron component (in particular, at optical and X-ray frequencies) of most blazars, including 1ES 1101-232 (e.g., Remillard et al. 1989; Romero et al. 1999; Wolter et al. 2000). We therefore argue that the rapidly variable synchrotron emission originates on substantially smaller (sub-pc) scales along the jet than the VHE emission.

3. Results for 1ES 1101-232

Based on the similarity of the optical and X-ray characteristics of 1ES 1101-232 (Remillard et al. 1989; Falomo et al. 1994) with previously known TeV blazars, Wolter et al. (2000) and Costamante & Ghisellini (2002) predicted that 1ES 1101-232 might be detectable by the now operating generation of VHE γ-ray detectors. Prompted by these predictions, Aharonian et al. (2006) performed three sets of coordinated optical – X-ray – VHE γ-ray observations in 2004 and 2005, which resulted in its detection at VHE γ-rays. This made 1ES 1101-232 the highest-redshift BL Lac object detected in VHE γ-rays at the time (now surpassed by 1ES 1011+496 at $z = 0.212$, Albert et al. (2007)) and the second-farthest VHE γ-ray source known, only topped by the quasar-type blazar 3C 279 at $z = 0.538$, for which the MAGIC collaboration recently claimed a VHE detection (Teshima et al. 2007). The H.E.S.S. observations of 1ES 1101-232 provided no evidence for variability at VHE γ-rays. Figs. 1 and 2 show the simultaneous optical – X-ray – VHE γ-ray spectra during two of the three H.E.S.S. observing windows, where the VHE γ-ray points have been corrected for $\gamma\gamma$ absorption by the IIBR as described in Aharonian et al. (2007a).

Unfortunately, all of the known observational facts on 1ES 1101-232 do not allow very



Fig. 1.— Simultaneous optical – X-ray – VHE γ-ray SED of 1ES 1101-232 on March 5 – 16, 2005 (from Aharonian et al. 2007a). The VHE measurements have been corrected for absorption by the IIBR as described in Aharonian et al. (2006). The optical data points are an upper and a lower R-band limit, respectively, derived from simultaneous ROTSE 3c observations in the 400 – 900 nm bandpass (Aharonian et al. 2007a). The curves indicate a model fit using an extended version of the code of Böttcher & Chiang (2002). The H.E.S.S. data have been fitted with $\Gamma = 15$, $B' = 10\,\mu$G, $\gamma'_1 = 100$, $\gamma'_2 = 6.0 \times 10^6$, $q = 1.5$, and $L_{\rm jet} = 1.5 \cdot 10^{41}$ ergs s^{-1}. The individual curves represent: dot-dashed = CMB Compton; double-dot-dashed = SSC; solid = total emission from the VHE emitting region. The dashed curve, fitting the synchrotron component, has been computed with parameters more appropriate to the inner-jet region, with $\Gamma = 25$, $\gamma'_1 = 1.3 \times 10^5$, $\gamma'_2 = 7.5 \times 10^5$, $q = 3.2$, $L_{\rm jet} = 2.0 \times 10^{40}$ ergs s^{-1}, $R'_B = 6 \cdot 10^{16}$ cm, and a magnetic field in equipartition with the relativistic electron plasma, at $B' = 0.06$ G. The X-ray/γ-ray portion of the dashed curve represents the inner-jet SSC emission.

meaningful constraints on the parameters of the kpc-scale jets of 1ES 1101-232 beyond a constraint on the overall jet power: The power injected into the jet on kpc scales obviously needs to exceed the observed radio power of the extended radio structure at 1.5 GHz, $P_{1.5}^{\rm ext} \approx 3.8 \times 10^{40}$ erg/s (Laurent-Muehleisen et al. 1993). Following arguments presented for the X-ray emission from extended jets of other AGN, indicating that the jets remain highly relativistic out to kpc scales (e.g. Tavecchio et al. 2000; Atoyan & Dermer 2004; Schwartz et al. 2006), we assume a characteristic Lorentz factor of $\Gamma = D = 15$. Based on the discussion in the previous section, an extension of the CMB Compton emission into the TeV regime requires electrons with energies up to $\gamma_2' \gtrsim 2.7 \cdot 10^6$. In order for the (non-variable) synchrotron emission from the TeV emission region not to dominate the observed X-ray emission, a magnetic field of $B' \lesssim 675\,\mu$G is required.

In order to evaluate the resulting broadband spectrum from a relativistic jet component, we have incorporated the process of CMB Compton scattering into the jet radiation transfer code described in detail in Böttcher & Chiang (2002). We used the head-on approximation for Compton scattering in the Thomson regime to treat the Compton scattering process and describe the CMB as a thermal blackbody at $T_{\rm CMB} = 2.72\,(1+z)$ K.

A reasonable representation of the VHE γ-ray spectrum as shown in Fig. 1 can be achieved using an electron injection index of $q = 1.5$, a luminosity injected in relativistic electrons into the kpc-scale jet of $L_{\rm jet} = 1.5 \times 10^{41}$ ergs s^{-1}, and a co-moving magnetic field of $B' = 10\,\mu$G. Obviously, these parameters are not tightly constrained, given that only the rather small spectral range of the VHE spectrum is fitted. The fits are insensitive to a low-energy cutoff ${\gamma'}_1$ of the electron spectrum, at least as long as ${\gamma'}_1 \lesssim 10^3$. The same choice of parameters provides an equally acceptable fit to the SED of June 5 – 10, 2004, as shown in Fig. 2.

As expected (and required), the predicted synchrotron emission from the extended jet region is about two orders of magnitude below the measured SED for both observation periods. In order to provide a fit to the synchrotron component of 1ES 1101-232, we chose parameters more appropriate for the inner (sub-pc) region of the jet, as listed in the figure captions. Note that the jet powers quoted there refer only to the power carried by the ultrarelativistic particle content. In the fits to the synchrotron component, we have chosen the magnetic field in equipartition with the ultrarelativistic electron population in the emission region in order to reduce the number of free parameters.

Although the parameters for the fits to both the synchrotron and the VHE components are not very well constrained, the general picture that emerges is that the CMB-Compton interpretation for the VHE emission requires a substantially larger kinetic luminosity of ultrarelativistic electron injection into the emission region on large scales than inferred for



Fig. 2.— Same as Fig. 1, but for the SED of June 5 – 10, 2004. Here, optical and X-ray data were obtained from simultaneous *XMM-Newton* observations (Aharonian et al. 2007a). Inner-jet parameters used for the synchrotron fit were: $\Gamma = 25$, $\gamma'_1 = 1.8 \times 10^5$, $\gamma'_2 = 1.5 \times 10^6$, $q = 2.5$, $L_{\rm jet} = 1.45 \times 10^{40}$ ergs s^{-1}, $R'_B = 6 \cdot 10^{16}$ cm, and $B' = 0.05$ G.

the synchrotron component at sub-pc scales. This suggests that a substantial fraction of the total jet luminosity is carried out to large distances by the non-leptonic content of the jet. If our assumption of a magnetic field in equipartition with relativistic electrons on sub-pc scales is realistic, this would suggest that most of the kinetic power of the jet might be carried by hadrons and dissipated on pc – kpc scales, leading to the acceleration of ultrarelativistic electrons in the extended jet. This is in perfect agreement with results of several other authors, suggesting that the pc-scale jets of blazars are dominated in number by electron-positron pairs, responsible for the high-energy emission produced at sup-pc scales, while the bulk of the kinetic power of the jets is carried by a relativistic hadron population (e.g. Sikora & Madejski 2000; Ghisellini & Celotti 2001; Kino & Takahara 2004; Sikora et al. 2005).

4. Summary and Discussion

We have presented a possible explanation for the hard VHE spectra of high-redshift γ-ray blazars, when corrected for the expected amount of $\gamma\gamma$ absorption by the IIBR. We suggest that ultrarelativistic electrons may be accelerated on large (kpc) scales, where magnetic fields are so low that synchrotron cooling becomes negligible compared to Compton cooling on the CMB. Consequently, VHE emission from Compton upscattering of the CMB by ultrarelativistic electrons may produce a separate, slowly variable VHE component beyond the rapidly variable broadband continuum from radio through GeV γ-ray energies that might be dominated by emission from the sub-pc scale, inner regions of the jet.

As of the time of writing of this manuscript, the VHE emission of most distant TeV blazars, e.g., 1ES 1101-232 (Aharonian et al. 2007a), 1ES 0229+200 (Aharonian et al. 2007c), 1ES 1011+496 (Albert et al. 2007), has not shown evidence for variability on any time scale, while the nearby TeV blazars show substantial variability on time scales from years down to a few minutes (e.g. Aharonian et al. 2007b). This could indicate that CMB Compton scattering constitutes an underlying non-varying VHE component in TeV blazars, which would be more likely to be detected in high-redshift sources since the energy density of the CMB increases with redshift as $(1+z)^4$. However, one has to caution that small-amplitude variability would generally be much harder to detect for the much fainter high-redshift TeV sources.

We have applied this idea to two simultaneous SEDs of 1ES 1101-232. Although the parameters of our fits are rather poorly constrained, they seem to indicate that only a small portion of the kinetic energy of the jet is dissipated on sub-pc scales, while the bulk of this energy is transported outward to kpc scales, in agreement with earlier suggestions that the kinetic power of blazar jets might generally be dominated by their hadronic content and is only dissipated on large scales.

Alternative explanations for the very hard VHE emissions include Compton upscattering of infrared emission from dust in the vicinity of the AGN (e.g., Błażejowski et al. 2000), or signatures of pion decay in hadronic blazar models (e.g., Mücke et al. 2003). However, it should be noted that Compton scattering of infrared radiation with a characteristic wavelength of $\lambda = \lambda_1\,\mu$m would also be affected by Klein-Nishina effects at energies of $E \gtrsim 0.2\,\lambda_1$ TeV, so a photon index of $\Gamma_{\rm int} \sim 1.5$ out to multi-TeV energies might be difficult to explain with this mechanism. It is also possible to produce VHE spectra of arbitrary hardness if they are affected by $\gamma\gamma$ absorption intrinsic to the source through interaction with narrow-band emission from the vicinity of the AGN (Aharonian et al. 2008). However, it is not obvious why such a feature should be observed preferentially in high-redshift TeV blazars, and why it would lead to less variable TeV emission than in the low-redshift TeV blazars, if the lack of variability is confirmed by future observations.

Our CMB-Compton interpretation for the VHE emission of 1ES 1101-232 is obviously not applicable to high-redshift VHE sources that seem to show substantial variability on time scales of days or even shorter, e.g., 3C279 (Teshima et al. 2007) since it requires a pc-scale extent of the emission region. The interpretation of the unexpectedly hard VHE spectrum through such a scenario would predict that any emission at lower (GeV) energies might be produced in the inner (sub-pc) region of the jet and be much more rapidly variable than the VHE emission. Whether this is the correct interpretation for the hard spectrum of 1ES 1101-232 or not, a Compton-scattered CMB component from blazar jet emission will accompany synchrotron emission and could potentially be observed with γ-ray telescopes.

We thank the referee for a very helpful and constructive report which helped to improve our manuscript substantially. The work of M. Böttcher as an ASEE Summer Research Faculty Fellowship and J. Finke as an NRC/NRL postdoctoral research associate at the Naval Research Laboratory was supported by NASA *GLAST* Science Investigation DPR-S-1563-Y. The work of C. D. is supported by the Office of Naval Research.

REFERENCES

Aharonian, F., et al., 2006, Nature, 440, 1018

Aharonian, F., et al., 2007a, A&A, 470, 475

Aharonian, F., et al., 2007b, ApJ, 664, L71

Aharonian, F., et al., 2007c, A&A, 475, L9

Aharonian, F., Khangulyan, D., & Costamante, L., 2008, MNRAS, submitted (astro-ph/0801.3198)

Albert, J., et al., 2007, ApJ, 667, L21

Atoyan, A., & Dermer, C. D. 2004, ApJ, 613, 151

Błażejowski, M., et al., 2000, ApJ, 545, 107

Böttcher, M., & Chiang, J., 2002, ApJ, 581, 127

Costamante, L., & Ghisellini, G., 2002, A&A, 384, 384, 56

Derishev, E. V., Aharonian, F. A., Kocharovsky, V. V., & Kocharovski, Vi. V., 2003, Phys. Rev. D, 68, 04003-1

Dermer, C. D. 2007, in 30th ICRC, Merida, Yucatan, Mexico, ArXiv e-prints, 711, arXiv:0711.2804

Dwek, E., & Krennrich, F., 2005, 618, 657

Falomo, R., Scarpa, R., & Bersanelli, M. 1994, ApJS, 93, 125

Finke, J. D., Dermer, C. D., & Böttcher, M., 2008, submitted

Ghisellini, G., & Celotti, A., 2001, MNRAS, 327, 739

Hauser, M. G., & Dwek, E., 2001, ARA&A, 39, 249

Katarzyński, K., et al., 2006, MNRAS, 368, L52

Kino, M., & Takahara, F., 2004, MNRAS, 349, 336

Laurent-Muehleisen, S. A., Kollgaard, R. I., Moellenbrock, G. A., & Feigelson, E. D., 1993, AJ, 106, 875

Madau, P., & Pozzetti, L., 2000, MNRAS, 312, L9

Mazin, D., & Raue, M., 2007, A&A, 471, 439

Mücke, A., Protheroe, R. J., Engel, R., Rachen, J. P., & Stanev, T., 2003, Astropart. Phys., 18, 593

Remillard, R. A., Tuohy, I. R., Brissenden, R. J. V., Buckley, D. A. H., Schwartz, D. A., Feigelson, E. D., & Tapia, S., 1989, ApJ, 345, 140

Rieger, F. M., & Duffy, P., 2006, ApJ, 652, 1044

Romero, G. E., Cellone, S. A., & Combi, J. A., 1999, ApJS, 135, 477

Sambruna, R. M., et al., 2004, ApJ, 608, 698

Schwartz, D. A., et al., 2006, ApJ, 640, 592

Sikora, M., & Madejski, G., 2000, ApJ, 534, 109

Sikora, M., et al., 2005, ApJ, 625, 72

Stawarz, L., & Ostrowski, M., 2002, ApJ, 578, 763

Stecker, F. W., Malkan, M. A., & Scully, S. T., 2006, ApJ, 648, 774 (Erratum: ApJ, 658, 1392 [2007])

Stecker, F. W., Baring, M. G., & Summerlin, E. J., 2007, ApJ, 667, L29

Stecker, F. W., & Scully, S. T., 2006, ApJ, 652, L9

Stern, B. E., & Poutanen, J., 2008, MNRAS, 383, 1695

Tavecchio, F., Maraschi, L, Sambruna, R. M., & Urry, C. M., 2000, ApJL, 544, L23

Teshima, M., et al. 2007, ArXiv e-prints, 709, arXiv:0709.1475

Virtanen, J. J. P., & Vainio, R., 2005, ApJ, 621, 313

Wolter, A., Tavecchio, F., Caccianiga, A., Ghisellini, G., & Tagliaferri, G., 2000, A&A, 357, 429

This preprint was prepared with the AAS LaTeX macros v5.2.